

Prenatal Joubert Syndrome and Related Disorders (JSRD) Panel Sequence Analysis and Exon-Level Deletion/Duplication Testing of 29 Genes

Panel Gene List

AHI1, ARL13B, B9D1, B9D2, C5orf42, CC2D2A, CEP41, CEP104, CEP120, CEP290, CSPP1, IFT172, INPP5E, KIAA0586, KIF7*, MKS1, NPHP1, NPHP3, OFD1, RPGRIP1L, TCTN1, TCTN2, TCTN3, TMEM67, TMEM138, TMEM216, TMEM231, TMEM237, TTC21B.

*Only whole gene deletions/duplications may be detected for the KIF7 gene

Clinical Features

Joubert syndrome is a brain malformation disorder characterized by the presence of a midbrain-hindbrain abnormality called the molar tooth sign, which results from cerebellar hypoplasia, thickened superior cerebellar peduncles, and a deepened interpeduncular fossa.¹⁻³ Affected individuals demonstrate hypotonia, developmental delay, and variable cognitive impairment. Breathing abnormalities and oculomotor apraxia may also be observed. Retinal dystrophy, renal disease, occipital encephalocele, polydactyly, hepatic fibrosis, and other abnormalities are seen in variant forms of this disorder. Renal disease can range from nephronophthisis to cystic renal dysplasia.⁴ Joubert syndrome shares phenotypic overlap with other ciliopathies, including Bardet-Biedl, COACH, and Meckel-Gruber syndromes, and all of these disorders show significant clinical variability.^{1,5-8} Meckel-Gruber syndrome, in particular, shows close overlap with Joubert syndrome and is characterized by occipital encephalocele, cystic kidneys, hepatic defects, and other anomalies.⁹ A diagnostic algorithm has been proposed to address the clinical and genetic heterogeneity and aid in the molecular diagnosis of Joubert syndrome.^{1,2}

Joubert syndrome and related disorders (JSRD) are identified in the prenatal period by the presence of the molar tooth sign on ultrasound.¹ Additional features including other structural brain malformations, encephalocele, renal disease, polydactyly, and cleft lip/palate may also be seen.¹ Other imaging methods such as 3D ultrasound or fetal MRI can be used to evaluate and diagnose JSRD in utero.¹ Due to genetic heterogeneity and overlapping phenotypes, the specific diagnosis cannot be determined accurately with imaging alone. When available, genetic testing can aid in determining the precise diagnosis after the differential has been established by imaging.

Inheritance Patterns/Genetics

The prevalence of Joubert syndrome and its variant forms is estimated to be approximately 1:100,000.¹ Many genes have been associated with Joubert syndrome, although some are very rarely involved.¹⁰ These genes have important functions in the development of cilia in various organs.³ Cilia play a role in intraflagellar transport, cell division, tissue differentiation, establishment of body axis, growth, and mechanosensation involved in cellular signaling processes.^{11,12} The differential diagnosis for Joubert syndrome includes other ciliopathies, such as Bardet-Biedl, COACH, and Meckel-Gruber syndromes.^{1,5-8} Meckel-Gruber syndrome is caused by variants in several of the Joubert syndrome-related genes.¹³ Renal disease is also a common feature between Joubert syndrome and nephronophthisis; most variants in the NPHP1 gene are associated with isolated nephronophthisis, but approximately 9% of individuals with NPHP1 deletions have features of Joubert syndrome.¹⁴ The central nervous system, renal epithelium, and photoreceptor cells are commonly affected in this group of disorders.

The GeneDx Joubert Syndrome Panel includes 29 genes that are involved in Joubert syndrome and related disorders (JSRD). Variants in these genes typically have a loss-of-function effect and include missense, nonsense, splicing, and insertion-deletion changes, as well as exonic deletions.

Test Methods

Using genomic DNA extracted from the submitted specimen, the complete coding regions and splice site junctions of the genes tested are enriched using a proprietary targeted capture system developed by GeneDx for next-generation sequencing with CNV calling (NGS-CNV). The enriched targets are simultaneously sequenced with paired-end reads on an Illumina platform. Bi-directional sequence reads are assembled and aligned to reference sequences based on NCBI RefSeq transcripts and human genome build GRCh37/UCSC hg19. After gene specific filtering, data are analyzed to identify sequence variants and most deletions and duplications involving coding exons; however, technical limitations and inherent sequence properties effectively reduce this resolution for some genes. Alternative sequencing or copy number detection methods are used to analyze or confirm regions with inadequate sequence or copy number data by NGS. Reportable variants include pathogenic variants, likely pathogenic variants and variants of uncertain significance. Likely benign and benign variants, if present, are not routinely reported but are available upon request.

The technical sensitivity of sequencing is estimated to be >99% at detecting single nucleotide events. It will not reliably detect deletions greater than 20 base pairs, insertions or rearrangements greater than 10 base pairs, or low-level mosaicism. The copy number assessment methods used with this test cannot reliably detect copy number variants of less than 500 base pairs or mosaicism and cannot identify balanced chromosome aberrations. Assessment of exon-level copy number events is dependent on the inherent sequence properties of the targeted regions, including shared homology and exon size. Gene specific exclusions for exon-level deletion/duplication testing for this panel are: KIF7 gene, only whole gene deletions or duplications may be detected.

Additionally, genotype analysis of maternal and fetal DNA for several polymorphic markers to test for maternal cell contamination will be performed. Therefore, in all prenatal cases, a maternal sample should accompany the fetal sample.

Test Sensitivity

The clinical sensitivity of sequencing and deletion/duplication analysis of the genes included in this panel in prenatal cases ascertained based on fetal ultrasound abnormalities is currently unknown, and the clinical sensitivity of this panel depends in part on the clinical phenotype of the fetus. Specific information about the diagnostic yield for each gene in selected populations is summarized in the following table(s).

Gene	Protein	Inheritance	Disease Associations
AHI1	Abelson helper integration site 1 (Joubertin)	AR	6-16% of JSRD ¹⁵⁻¹⁹
ARL13B	ADP-ribosylation factor-like 13B	AR	Rare in JSRD ^{20,21}
B9D1	B9 domain-containing protein 1	AR	Rare in JSRD ^{17,22,23}
B9D2	B9 domain-containing protein 2	AR	Rare in JSRD ^{28,87}
C5orf42	Chromosome 5 open reading frame 42	AR	7-25% of JSRD ^{15-17,24} ; 45% of JSRD in French-Canadian ²⁵ ; 82% of Oral-facial-digital syndrome type 6 (OFDVI) ²⁶
CC2D2A (MKS6)	Coiled-coil and C2 domainscontaining protein 2A	AR	2-10% of JSRD ^{15,17,27-33}
CEP41	Centrosomal protein, 41kDa	AR	Rare in JSRD ³⁴

CEP104	Centrosomal protein, 104kDa	AR	Rare in JSRD ⁸⁸
CEP120	Centrosomal protein, 120kDa	AR	Rare in JSRD ^{89,90}
CEP290 (NPHP6)	Centrosomal protein, 290kDa	AR	2-25% of JSRD ^{15-17,32,33,35-40} ; 50% in JSRD with cerebello-oculo-renal phenotype (CORS) ⁴¹
CSPP1	Centrosome and spindle pole associated protein 1	AR	2-5% of JSRD ^{15,28,42-44}
IFT172	Intraflagellar transport 172	AR	Rare in JSRD ^{28,45,46}
INPP5E	Inositol polyphosphate-5phosphatase, 72kDa	AR	2-4% of JSRD ^{15,17,28,47,48}
KIAA0586	TALPID3 protein	AR	2-7% of JSRD ⁹¹⁻⁹⁵
KIF7*	Kinesin family member 7	AR	Rare in JSRD, fetal hydroletharus and acrocallosal syndromes ^{15,28,49-52}
MKS1	MKS1 B9-domain containing protein (Meckel syndrome type 1 protein)	AR	1-2% of JSRD ^{15,23,28} ; 7-30% of Meckel Gruber syndrome ^{9,13,33,53}
NPHP1	Nephrocystin 1	AR	2-7% of JSRD (homozygous gene deletion) ^{14,54-56}
NPHP3	Nephrocystin 3	AR	3% of JSRD ^{36,57-60}
OFD1 (CXorf5)	Oral-facial-digital syndrome 1 protein	XL	Rare in JSRD ^{15,17,28,61-64}
RPGRIP1L (NPHP8)	RPGRIP1-like protein	AR	2-5% in Joubert syndrome (mostly cerebro-renal type) ^{15,17,28,29,33,65-68}
TCTN1	Tectonic family member 1	AR	Rare in JSRD ^{15,16,69,70}
TCTN2	Tectonic family member 2	AR	Rare in JSRD ^{15,28,64,71-73}
TCTN3	Tectonic family member 3	AR	Rare in JSRD ^{15,71,74}
TMEM67	Transmembrane protein 67	AR	1-10% of JSRD ^{6,15,17,28,32,36,75,76} ; 15-28% of Meckel
(MKS3)	(Meckelin)		Gruber syndrome ^{9,33} ; 75-83% of JSRD with liver involvement (including COACH syndrome) ²⁹
TMEM138	Transmembrane protein 138	AR	Rare in JSRD ^{15,33,77}
TMEM216	Transmembrane protein 216	AR	3% of JSRD ^{28,77,78} ; Up to 100% of JSRD in Ashkenazi Jewish ^{78,79} ; Rare in Oral-facial-digital syndrome type 6 (OFDVI) ²⁶
TMEM231	Transmembrane protein 231	AR	Rare in Joubert and Meckel Gruber syndromes ^{17,80-82}
TMEM237 (ALS2CR4)	Transmembrane protein 237	AR	Rare in Joubert and Meckel Gruber syndromes ^{15,16,33,53,83}
TTC21B	Tetratricopeptide repeat domaincontaining protein 21B	AR	Unknown in JSRD ^{32,84-86}

* Only whole gene deletions/duplications may be detected for the KIF7 gene

REFERENCES:

1. Parisi M, Glass I. Joubert Syndrome and Related Disorders. 2003 Jul 9 [Updated 2013 Apr 11]. In: Pagon RA, Adam MP, Bird TD, et al., editors. GeneReviews™ [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2013. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK1325/>;
2. Parisi et al. (2009) American Journal of Medical Genetics. Part C, Seminars In Medical Genetics 151C (4):326-40 (PMID: 19876931)
3. Sattar et al. (2011) Developmental Medicine And Child Neurology 53 (9):793-8 (PMID: 21679365)
4. Doherty et al. (2009) Seminars In Pediatric Neurology 16 (3):143-54 (PMID: 19778711)
5. Abdelhamed et al. (2013) Human Molecular Genetics 22 (7):1358-72 (PMID: 23283079)
6. Baala et al. (2007) American Journal Of Human Genetics 80 (1):186-94 (PMID: 17160906)
7. Badano et al. (2006) Annual Review Of Genomics And Human Genetics 7 :125-48 (PMID: 16722803)
8. Myktyyn et al. (2007) Nature Genetics 39 (7):818-9 (PMID: 17597771)
9. Khaddour et al. (2007) Human Mutation 28 (5):523-4 (PMID: 17397051)
10. Online Mendelian Inheritance in Man; <http://omim.org>
11. Goetz et al. (2010) Nat. Rev. Genet. 11 (5):331-44 (PMID: 20395968)
12. Lee et al. (2010) Neurobiology Of Disease 38 (2):167-72 (PMID: 20097287)
13. Chen et al. (2007) Taiwanese Journal Of Obstetrics & Gynecology 46 (1):9-14 (PMID: 17389183)
14. Caridi et al. (2006) Kidney International 70 (7):1342-7 (PMID: 16900087)
15. Akizu et al. (2014) American Journal Of Human Genetics 94 (1):80-6 (PMID: 24360807)
16. Alazami et al. (2012) Human Mutation 33 (10):1423-8 (PMID: 22693042)
17. Kroes et al. (2015) European Journal Of Human Genetics : Ejhg : (PMID: 25920555)
18. Parisi et al. (2006) Journal Of Medical Genetics 43 (4):334-9 (PMID: 16155189)
19. Valente et al. (2006) Annals Of Neurology 59 (3):527-34 (PMID: 16453322)
20. Cantagrel et al. (2008) American Journal Of Human Genetics 83 (2):170-9 (PMID: 18674751)
21. Thomas et al. (2015) Eur. J. Hum. Genet. 23 (5):621-7 (PMID: 25138100)
22. Hopp et al. (2011) Human Molecular Genetics 20 (13):2524-34 (PMID: 21493627)
23. Romani et al. (2014) Orphanet Journal Of Rare Diseases 9 :72 (PMID: 24886560)
24. Romani et al. (2015) Human Genetics 134 (1):123-6 (PMID: 25407461)
25. Srour et al. (2012) American Journal Of Human Genetics 90 (4):693-700 (PMID: 22425360)
26. Lopez et al. (2014) Human Genetics 133 (3):367-77 (PMID: 24178751)
27. Bachmann-Gagescu et al. (2012) Journal Of Medical Genetics 49 (2):126-37 (PMID: 22241855)
28. Bachmann-Gagescu et al. (2015) J. Med. Genet. 52(8):514-22 (PMID: 26092869)
29. Doherty et al. (2010) Journal Of Medical Genetics 47 (1):8-21 (PMID: 19574260)
30. Gordon et al. (2008) American Journal Of Human Genetics 83 (5):559-71 (PMID: 18950740)
31. Mougou-Zerelli et al. (2009) Human Mutation 30 (11):1574-82 (PMID: 19777577)
32. Otto et al. (2011) Journal Of Medical Genetics 48 (2):105-16 (PMID: 21068128)
33. Szymanska et al. (2012) Cilia 1 (1):18 (PMID: 23351400)
34. Lee et al. (2012) Nature Genetics 44 (2):193-9 (PMID: 22246503)
35. Brancati et al. (2007) American Journal Of Human Genetics 81 (1):104-13 (PMID: 17564967)
36. Halbritter et al. (2012) Journal of Medical Genetics 49 (12):756-67 (PMID: 23188109).
37. Helou et al. (2007) J. Med. Genet. 44 (10):657-63 (PMID: 17617513)
38. Sayer et al. (2006) Nature Genetics 38 (6):674-81 (PMID: 16682973)
39. Travaglini et al. (2009) American Journal of Medical Genetics. Part A 149A (10):2173-80 (PMID: 19764032)
40. Valente et al. (2006) Nature Genetics 38 (6):623-5 (PMID: 16682970)
41. Coppeters et al. (2010) Human Mutation 31 (10):1097-108 (PMID: 20690115)
42. Ben-Omran et al. (2015) Am. J. Med. Genet. A 167 (10):2478-80 (PMID: 25997910)
43. Shaheen et al. (2014) American Journal Of Human Genetics 94 (1):73-9 (PMID: 24360803)
44. Tuz et al. (2014) American Journal Of Human Genetics 94 (1):62-72 (PMID: 24360808)
45. Bujakowska et al. (2015) Hum. Mol. Genet. 24 (1):230-42 (PMID: 25168386)
46. Halbritter et al. (2013) American Journal Of Human Genetics 93 (5):915-25 (PMID: 24140113)
47. Bielas et al. (2009) Nature Genetics 41 (9):1032-6 (PMID: 19668216)
48. Travaglini et al. (2013) European Journal Of Human Genetics : Ejhg 21 (10):1074-8 (PMID: 23386033)
49. Dafinger et al. (2011) The Journal Of Clinical Investigation 121 (7):2662-7 (PMID: 21633164)
50. Putoux et al. (2011) Nature Genetics 43 (6):601-6 (PMID: 21552264)
51. Putoux et al. (2012) Journal Of Medical Genetics 49 (11):713-20 (PMID: 23125460)
52. Walsh et al. (2013) European Journal Of Medical Genetics 56 (1):39-42 (PMID: 23142271).
53. Shaheen et al. (2013) European Journal Of Human Genetics : Ejhg 21 (7):762-8 (PMID: 23169490)
54. Castori et al. (2005) Journal Of Medical Genetics 42 (2):e9 (PMID: 15689444)
55. Parisi et al. (2004) American Journal Of Human Genetics 75 (1):82-91 (PMID: 15138899)
56. Tory et al. (2007) Journal Of The American Society Of Nephrology : Jasn 18 (5):1566-75 (PMID: 17409309)
57. Bergmann et al. (2008) American Journal Of Human Genetics 82 (4):959-70 (PMID: 18371931)
58. Calinescu-Tuleasca et al. (2013) Eur. J. Pediatr. 172 (7):877-81 (PMID: 21845392)
59. Copelovitch et al. (2013) Am. J. Med. Genet. A 161A (7):1743-9 (PMID: 23686967)
60. Leeman et al. (2014) J Perinatol 34 (5):410-1 (PMID: 24776604)
61. Bisschoff et al. (2013) Hum. Mutat. 34 (1):237-47 (PMID: 23033313)
62. Darmency-Stamboul et al. (2013) Eur J Med Genet 56 (6):301-8 (PMID: 23523602)
63. Field et al. (2012) European Journal Of Human Genetics : Ejhg 20 (7):806-9 (PMID: 22353940)
64. Juric-Sekhar et al. (2012) Acta Neuropathol. 123 (5):695-709 (PMID: 22331178)
65. Arts et al. (2007) Nature Genetics 39 (7):882-8 (PMID: 17558407)
66. Brancati et al. (2008) Clinical Genetics 74 (2):164-70 (PMID: 18565097)
67. Delous et al. (2007) Nature Genetics 39 (7):875-81 (PMID: 17558409)
68. Wolf et al. (2007) Kidney International 72 (12):1520-6 (PMID: 17960139)

- 69. Garcia-Gonzalo et al. (2011) Nature Genetics 43 (8):776-84 (PMID: 21725307)
- 70. Reiter et al. (2006) Genes Dev. 20 (1):22-7 (PMID: 16357211)
- 71. Huppke et al. (2015) European Journal Of Human Genetics : Ejhg 23 (5):616-20 (PMID: 25118024)
- 72. Sang et al. (2011) Cell 145 (4):513-28 (PMID: 21565611)
- 73. Shaheen et al. (2011) Human Mutation 32 (6):573-8 (PMID: 21462283)
- 74. Thomas et al. (2012) American Journal Of Human Genetics 91 (2):372-8 (PMID: 22883145)
- 75. Brancati et al. (2009) Human Mutation 30 (2):E432-42 (PMID: 19058225)
- 76. Iannicelli et al. (2010) Human Mutation 31 (5):E1319-31 (PMID: 20232449)
- 77. Lee et al. (2012) Science (New York, N.Y.) 335 (6071):966-9 (PMID:22282472)
- 78. Valente et al. (2010) Nature Genetics 42 (7):619-25 (PMID: 20512146)
- 79. Edvardson et al. (2010) American Journal Of Human Genetics 86 (1):93-7 (PMID: 20036350)
- 80. Srour et al. (2012) Journal Of Medical Genetics 49 (10):636-41 (PMID: 23012439)
- 81. Roberson et al. (2015) J. Cell Biol. 209 (1):129-42 (PMID: 25869670)
- 82. Shaheen et al. (2013) Journal Of Medical Genetics 50 (3):160-2 (PMID: 23349226)
- 83. Huang et al. (2011) American Journal Of Human Genetics 89 (6):713-30 (PMID: 22152675)
- 84. Davis et al. (2011) Nature Genetics 43 (3):189-96 (PMID: 21258341)
- 85. Halbritter et al. (2013) Human Genetics 132 (8):865-84 (PMID: 23559409)
- 86. Redin et al. (2012) Journal of Medical Genetics 49 (8):502-12 (PMID: 22773737)
- 87. Dowdle et al. (2011) American Journal Of Human Genetics 89 (1):94-110 (PMID: 21763481)
- 88. Srour et al. (2015) Am. J. Hum. Genet. 97 (5):744-53 (PMID: 26477546)
- 89. Shaheen et al. (2015) Hum. Mol. Genet. 24 (5):1410-9 (PMID: 25361962)
- 90. Roosing et al. (2016) J. Med. Genet. 53 (9):608-15 (PMID: 27208211)
- 91. Alby et al. (2015) Am. J. Hum. Genet. 97 (2):311-8 (PMID: 26166481)
- 92. Bachmann-Gagescu et al. (2015) Hum. Mutat. 36 (9):831-5 (PMID: 26096313)
- 93. Malicdan et al. (2015) J. Med. Genet. 52 (12):830-9 (PMID: 26386044)
- 94. Roosing et al. (2015) Elife 4 :e06602 (PMID: 26026149) 95. Vilboux et al. (2017) Genet. Med. : (PMID: 28125082)